Grass and Forage Science

EDITED BY A. YOUNGER

VOLUME 50, 1995

The Journal of the British Grassland Society
published by Blackwell Science
Oxford London Edinburgh Boston
Melbourne Berlin Paris Vienna

© 1995 Blackwell Science
Osney Mead, Oxford OX2 0EL
25 John Street, London WCIN 2BL
23 Ainslie Place, Edinburgh EH3 6AJ
238 Main Street, Boston, MA
02142, USA
54 University Street, Carlton, Victoria 3054,
Australia

Other editorial officers

Blackwell Wissenschafts-Verlag GmbH,
Kurfürstendamm 57, 10707
Berlin, Germany
Arnette Blackwell SA, 224 Boulevard St Germain,
75007 Paris, France
Blackwell Wissenschafts-Verlag GmbH, Zehetnergasse 6,
A-1140 Wien, Austria

Authorization to photocopy items for internal or personal use, or the internal or personal use of specific clients, is granted by Blackwell Science for libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that the base fee of $11.50 per copy is paid directly to CCC, 222 Rosewood Drive, Suite 910, Danvers, MA 01923, USA.Special requests should be addressed to the Editor. 0142-5242/94 $11.50

ISSN 0142-5242

Printed in Great Britain by Whitstable Litho, Whitstable, UK

Contents

NUMBER 1, MARCH 1995

M.J. HILL, M.J. HOCKNEY, C.A. MULCAHY and G.RAPP. The effect of hydrogen cyanide potential (HCNp) and sward morphology on the relative acceptability to sheep of white clover and Caucasian clover herbage 1

D.R.EVANS, T.A. WILLIAMS, S. JONES and S.A. EVANS. The effect of blending white clover varieties and their contribution to a mixed grass/clover sward under continuous sheep stocking 10

W.J. FULKERSON and K.SLACK. Leaf number as a criterion for determining defoliation time for *Lolium perenne*: 2. Effect of defoliation frequency and height 16

J.P. FROST, R.POOTS, A KNIGHT, F.J. GORDON and F.N.J. LONG. Effect of forage matting rate on grass drying, rate of silage fermentation, silage intake and silage digestibility 21

R.J. ORR, P.D. PENNING, A.J. PARSONS and R.A. CHAMPION. Herbage intake and N excretion by sheep grazing monocultures or a mixture of grass and white clover 31

C. ÖSTLING and S. LINDGREN. Influences of enterobacteria on the fermentation and aerobic stability of grass silages 41

A.S. LAIDLAW, J.A. WITHERS and L.G. TOAL. The effect of surface height of swards continuously stocked with cattle on herbage production and clover content over four years 48

V.R. KANNEGANTI and S.R. KAFFKA. Forage availability from a temperate pasture managed with intensive rotational grazing 55

P.M. HAIGH. The composition of first-cut grass for ensilage in England and Wales from 1988 to 1991 (Research Note) 63

R.G. SIMONS. A non-destructive method for estimating lucerne shoot weights (Research Note) 68

J. WAYMAN, D.G. CHAMBERLAIN and A.G. WILLIAMS. A comparison of the ensilage, in an anaerobic cabinet, of grass either fresh or when thawed after deep freezing (Research Note) 71

C. RAVEL, G. CHARMET and F. BALFOURIER. Influence of the fungal endophyte *Acremonium lolii* on agronomic traits of perennial ryegrass in France (Research Note) 75

M. MERCHANT. The effect of pattern and severity of cutting on the vigour of the soft rush (*Juncus effusus* L.) (Research Note) 81

A. CHERRILL. Infestation of improved grasslands by *Juncus effusus* L. in the catchment of the River Tyne, Northern England: a field survey (Research Note) 85

NUMBER 2, JUNE 1995

S.V. PASUMARTY, S. HIGUCHI, T. MURATA and T. MATSUMARA. Influence of seed quality on seedling growth of white clover (*Trifolium repens* L.) 93

G.T. BARTHRAM and S.A. GRANT Interactions between variety and the timing of conservation cuts on species balance in *Lolium Perenne-Trifolium repens* swards 98

C.D. STUDDY, R.M. MORRIS and I. RIDGE. The effects of separated cow slurry liquor on soil and herbage nitrogen in *Phalaris arundinacea* and *Lolium perenne* 106

R.J. MERRY, M.S. DHANOA and M.K. THEODOROU. Use of freshly cultured lactic acid bacteria as silage inoculants 112

K.A.SMITH, D.R. JACKSON, R.J. UNWIN, G. BAILEY and I.HODGSON. Negative effects of winter- and spring-applied cattle slurry on the yield of herbage at simulated early grazing and first-cut silage 124

A.G. WILLIAMS, D.L. CRITTEN and A.M. REYNOLDS. A mathematical model of the aerobic deterioration of silage 132

S.V. PASUMARGTY, S. HIGUCHI, T. MURATA and T. MATSUMARA. Seed production in white clover (*Trifolium repens* L.). II. Effect of autumn defoliation on potential seed yield and seed yield components 147

P.W. WILKINS. Independence of dry matter yield and leaf yield among perennial ryegrass varieties differing in seasonal yield distribution 155

A.FISHER and D. WILMAN. Effect of interval between harvests and spring-applied fertilizer N on the growth of white clover in a mixed sward 162

R.F. WELLER, A. COOPER and M.S. DHANOA. The selection of winter wheat varieties for whole-crop cereal conservation (Research Note) 172

C. MATTHEW, A. CRESSWELL and R.J. HAGGAR. Characteristics of microsites with and without white clover in two field swards (Research Note) 178

NUMBER 3, SEPTEMBER 1995

W.M. MURPHY, A.D. MENA BARRETO and J.P.SILMAN. Sward dynamics of a smooth-stalked meadowgrass-dominant white clover sward rotationally grazed by cattle and/or sheep 183

W.M. MURPHY, A.D. MENA BARRETO, J.P. SILMAN and D.L. DINDAL. Cattle and sheep grazing effects on soil organisms, fertility and compaction in a smooth-stalked meadowgrass-dominant white clover sward 191

E.PIANO and P. ANNICCHIARICO. Persistence of Ladino white clover ecotypes and its relationship with other agronomic traits 195

R.J. WILKINS, M.J. GIBB and C.A. HUCKLE. Lactation performance of spring-calving dairy cows grazing mixed perennial ryegrass/white clover swards of differing composition and height 199

D.P. HANDA, P.R. SREENATH and S.K. RAJPALI. Uniformity trial with lucerne grown for fodder 209

T.W.J. KEADY and R.W.J. STEEN. The effects of treating low dry-matter, low digestibility grass with a bacterial inoculant on the intake and performance of beef cattle, and studies on its mode of action 217

D.A. DAVIES, M.FOTHERGILL, G.J. DANIEL and C.T. MORGAN. Animal production evaluation of herbage varieties. 2. Comparison of Aberystwyth S184, AberEndura and Grasslands Huia white clovers 227

L. HAZARD and M. GHESQUIÈRE. Evidence from the use of isozyme markers of competition in swards between short-leaved and long-leaved perennial ryegrass 241

R.F. CUSSEN, R.J. MERRY, A.P. WILLIAMS and J.K.S. TWEED. The effect of additives on the ensilage of forage of differing perennial ryegrass and white clover content 249

L. THOMPSON. Sites of photoperception in white clover 259

P.H. WILLIAMS and R.J. HAYNES. Effect of sheep, deer and cattle dung on herbage production and soil nutrient content 263

R.C. BINNIE and J.P. FROST. Some effects of applying undiluted silage effluent to grassland 272

P. DILLON, S. CROSSE, G. STAKELUM and F. FLYNN. The effect of calving date and stocking rate on the performance of spring-calving dairy cows 286

N.J. HUTCHINGS and I.S. KRISTENSEN. Modelling mineral nitrogen accumulation in grazed pasture: will more nitrogen leach from fertilized grass than unfertilized grass than unfertilized grass/clover? 300

NUMBER 4, DECEMBER 1995

G.E.J. FISHER and A.M. DOWDESWELL. The performance of

summer-calving cows grazing perennial ryegrass swards 315

A.H. MARSHALL. Peduncle characteristics, inflorescence survival and reproductive growth of white clover (*Trifolium repens* L.) 324

G.D. RUXTON and G.J. GIBSON. A mathematical model of aerobic deterioration of big-bale silage and its implications for the growth of Listeria monocytogenes 331

J.D. PATTERSON, A.S. LAIDLAW and J. MCBRIDE. The influence of autumn management and companion grass on the development of white clover over winter in mixed swards 345

K.F. EDGAR, D.A. KLESSA, J. FRAME and R.D. HARKESS. The nitrogen value of anaerobically digested sewage sludge applied to grassland during winter 353

F.W. KIRKHAM and J.R.B. TALLOWIN. The influence of cutting date and previous fertilizer treatment on the productivity and botanical composition of species-rich hay meadows on the Somerset Levels 365

G.R. EDWARDS, A.J. PARSONS, P.D. PENNING and J.A. NEWMAN. Relationship between vegetation state and bite dimensions of sheep grazing contrasting plant species and its implications for intake rate and diet selection 378

R.H. ARMSTRONG, E. ROBERTSON and E.A. HUNTER. The effect of sward height and its direction of change on the herbage intake, diet selection and performance of weaned lambs grazing ryegrass swards 389

W.J. HAMILTON, A.M. SIBBALD and D. FEIST. The effect of sward height on the liveweight gain of farmed yearling red deer (*Cervus elaphus*) stags 399

P. ANNICCHIARICO, F.BOZZO, G. PARENTE, F. GUSMEROLI, V. MAIR, O. MARGUERETTAZ and D. ORLANDI. Analysis of adaptation of grass/legume mixtures to Italian alpine and subalpine zones through an additive main effects and multiplicative interaction model 405

M.J. HAYES and E.D. WILLIAMS. Effects of timing and intensity of defoliation on growth and development of white clover seedlings 414

G.E.J. FISHER, D.J. ROBERTS and A.M. DOWDESWELL. The manipulation of grass swards for summer-calving dairy cows 424

M.J. BULLARD and T.J. CRAWFORD. Productivity of *Lotus corniculatus* L. (bird's-foot trefoil) in the UK

when grown under low-input conditions as spaced plants, monoculture swards or mixed swards 439

I. CALVIÈRE and M.DURU. Leaf appearance and senescence patterns of some pasture species 447

W.M. MURPHY, J.P. SILMAN and A.D. MENA BARRETO. A comparison of quadrat, capacitance meter, HFRO sward stick, and rising plate for estimating herbage mass in a smooth-stalked, meadowgrass-dominant white clover sward 452

I. PAPASTYLIANOU. Effect of rainfall and temperature on yield of *Vicia sativa* under rainfed Mediterranean conditions 456

G.E.J. FISHER and D.J. ROBERTS. The effects of stocking rate in spring, and topping on sward characteristics and dairy cow performance in poor grass growing season 461

Author Index 466